

УДК 517.518.23+517.548.2  
DOI 10.46698/w5793-5981-8894-o

ON POLETSKY-TYPE MODULUS INEQUALITIES  
FOR SOME CLASSES OF MAPPINGS<sup>#</sup>

S. K. Vodopyanov<sup>1</sup>

<sup>1</sup> Sobolev Institute of Mathematics,  
4 Akademika Koptyuga Ave., Novosibirsk 630090, Russia  
E-mail: vodopis@math.nsc.ru

**Abstract.** It is well-known that the theory of mappings with bounded distortion was laid by Yu. G. Reshetnyak in 60-th of the last century [1]. In papers [2, 3], there was introduced the two-index scale of mappings with weighted bounded  $(q, p)$ -distortion. This scale of mappings includes, in particular, mappings with bounded distortion mentioned above (under  $q = p = n$  and the trivial weight function). In paper [4], for the two-index scale of mappings with weighted bounded  $(q, p)$ -distortion, the Poletsky-type modulus inequality was proved under minimal regularity; many examples of mappings were given to which the results of [4] can be applied. In this paper we show how to apply results of [4] to one such class. Another goal of this paper is to exhibit a new class of mappings in which Poletsky-type modulus inequalities is valid. To this end, for  $n = 2$ , we extend the validity of the assertions in [4] to the limiting exponents of summability:  $1 < q \leq p \leq \infty$ . This generalization contains, as a special case, the results of recently published papers. As a consequence of our results, we also obtain estimates for the change in capacity of condensers.

**Key words:** quasiconformal analysis, Sobolev space, modulus of a family of curves, modulus estimate.

**AMS Subject Classification:** 30C65 (26B35, 31B15, 46E35).

**For citation:** Vodopyanov, S. K. On Poletsky-type Modulus Inequalities for Some Classes of Mappings, *Vladikavkaz Math. J.*, 2022, vol. 24, no. 4, pp. 58–69. DOI: 10.46698/w5793-5981-8894-o.

## 1. Introduction

The goal of this work is to show the application of results of [4] for output of Poletsky-type modulus inequalities for some classes of mappings. For doing this we formulate first the main result of [4], and then we provide how it can be applied for some concrete classes of mappings.

The main classes of mappings studied in [4] were defined in [2, 3].

**DEFINITION 1.** Let  $\omega: \mathbb{R}^n \rightarrow [0, \infty]$  be a measurable function, called a *weight*, with  $0 < \omega < \infty$  holding  $\mathcal{H}^n$ -almost everywhere, and  $\Omega \subset \mathbb{R}^n$  is a domain in  $\mathbb{R}^n$ . A mapping  $f: \Omega \rightarrow \mathbb{R}^n$  with  $n \geq 2$  is called a *mapping with (inner) bounded  $\omega$ -weighted  $(q, p)$ -codistortion*, or briefly,  $f \in \mathcal{I}\mathcal{D}(\Omega; q, p; \omega, 1)$ , where  $n - 1 \leq q \leq p < \infty$ , whenever

- (1)  $f$  is continuous, open and discrete;
- (2)  $f$  belongs to the Sobolev class  $W_{n-1, \text{loc}}^1(\Omega)$ ;
- (3) the Jacobian determinant satisfies  $\det Df(x) \geq 0$  for almost all  $x \in \Omega$ ;

---

<sup>#</sup> The study was carried out within the framework of the State contract of the Sobolev Institute of Mathematics, project № FWNF-2022-0006.

(4) the mapping  $f$  has *bounded codistortion*:  $\text{adj } Df(x) = 0$  a. e. on the set  $Z = \{x \in \Omega : \det Df(x) = 0\}$ ;

(5) the *local  $\omega$ -weighted  $(q, p)$ -codistortion function*

$$\Omega \ni x \mapsto \mathcal{K}_{q,p}^{\omega,1}(x, f) = \begin{cases} \frac{\omega^{\frac{n-1}{q}}(x) |\text{adj } Df(x)|}{\det Df(x)^{\frac{n-1}{p}}} & \text{if } \det Df(x) \neq 0, \\ 0 & \text{otherwise,} \end{cases} \quad (1)$$

belongs to  $L_{\varrho}(\Omega)$ , where  $\varrho$  satisfies  $\frac{1}{\varrho} = \frac{n-1}{q} - \frac{n-1}{p}$ , while  $\varrho = \infty$  for  $q = p$ .

Put  $\mathcal{K}_{q,p}^{\omega,1}(f; \Omega) = \|\mathcal{K}_{q,p}^{\omega,1}(\cdot, f) | L_{\varrho}(\Omega)\|$ .

DEFINITION 2. Let  $\omega: \mathbb{R}^n \rightarrow [0, \infty]$  be a measurable function, called a *weight*, with  $0 < \omega < \infty$  holding  $\mathcal{H}^n$ -almost everywhere, and  $\Omega \subset \mathbb{R}^n$  is a domain in  $\mathbb{R}^n$ . A mapping  $f: \Omega \rightarrow \mathbb{R}^n$  with  $n \geq 2$  is called a *mapping with (outer) bounded  $\omega$ -weighted  $(q, p)$ -distortion*, or briefly  $f \in \mathcal{O}\mathcal{D}(\Omega; q, p; \omega, 1)$ , with  $n - 1 \leq q \leq p < \infty$ , whenever:

- (1)  $f$  is continuous, open and discrete;
- (2)  $f$  belongs to the Sobolev class  $W_{n-1, \text{loc}}^1(\Omega)$ ;
- (3) the Jacobian determinant satisfies  $\det Df(x) \geq 0$  for a. e.  $x \in \Omega$ ;
- (4) the mapping  $f$  has *bounded distortion*:  $Df(x) = 0$  a. e. on the set  $Z = \{x \in \Omega : \det Df(x) = 0\}$ ;

(5) the *local  $\omega$ -weighted  $(q, p)$ -distortion function*

$$\Omega \ni x \mapsto K_{q,p}^{\omega,1}(x, f) = \begin{cases} \frac{\omega^{\frac{1}{q}}(x) |Df(x)|}{\det Df(x)^{\frac{1}{p}}} & \text{if } \det Df(x) \neq 0, \\ 0 & \text{otherwise,} \end{cases} \quad (2)$$

belongs to  $L_{\varkappa}(\Omega)$ , where  $\varkappa$  satisfies  $\frac{1}{\varkappa} = \frac{1}{q} - \frac{1}{p}$ , while  $\varkappa = \infty$  for  $q = p$ .

Put  $K_{q,p}^{\omega,1}(f; \Omega) = \|K_{q,p}^{\omega,1}(\cdot, f) | L_{\varkappa}(\Omega)\|$ .

REMARK 1. It is established in [3] that

$$\mathcal{O}\mathcal{D}(\Omega; q, p; \omega, 1) \subset \mathcal{I}\mathcal{D}(\Omega; q, p; \omega, 1) \quad (3)$$

in case of  $n - 1 < q \leq p < \infty$ .

For justifying (3) we refer to [3, Theorem 8] where it is proved that every mapping  $f: \Omega \rightarrow \Omega'$  of  $\mathcal{O}\mathcal{D}(\Omega; q, p; \omega, 1)$ ,  $n - 1 < q \leq p < \infty$ , belongs also to the class  $\mathcal{I}\mathcal{D}(\Omega; q, p; \omega, 1)$ , and the estimate

$$\|\mathcal{K}_{q,p}^{\omega,1}(\cdot, f) | L_{\varrho}(\Omega)\| \leq \|K_{q,p}^{\omega,1}(\cdot, f) | L_{\varkappa}(\Omega)\|^{n-1} \quad (4)$$

holds. (Here  $\varrho$  and  $\varkappa$  are defined after formulas (1) and (2) respectively).

In [4] it was proved the following result.

**Theorem 1** [4, Theorem 4.1]. *Let  $n - 1 < q \leq p < \infty$ . Suppose that  $f: \Omega \rightarrow \mathbb{R}^n$  is a mapping with with inner bounded  $\omega$ -weighted  $(q, p)$ -codistortion ( $f \in \mathcal{I}\mathcal{D}(\Omega; q, p; \omega, 1)$ ), while the weight function  $\theta(x) = \omega^{-\frac{n-1}{q-(n-1)}}(x)$  is locally summable. If  $\Gamma$  is a family of curves in the domain  $\Omega$  then we have the inequality*

$$(\text{mod}_s f(\Gamma))^{1/s} \leq \mathcal{K}_{q,p}^{\omega,1}(f; \Omega) (\text{mod}_r^{\theta} \Gamma)^{1/r}, \quad (5)$$

with  $s = \frac{p}{p-(n-1)}$  and  $r = \frac{q}{q-(n-1)}$ .

Below we recall the concept of the modulus of a family of curves (see [4] for more details).

A *curve* in  $\mathbb{R}^n$  is a continuous mapping  $\alpha: I \rightarrow \mathbb{R}^n$ , where  $I$  is an interval in  $\mathbb{R}$ , that is, a set of the form  $\langle a, b \rangle$ , where each angular parenthesis can be either round or square,  $a, b \in \mathbb{R}$  with  $a \leq b$ . We also allow infinite intervals. A curve  $\alpha$  is called closed (open) if the interval  $I$  is compact (open). Put  $|\alpha| = \alpha(I)$ . The expression  $\gamma' \subset \gamma$  will mean that the curve  $\gamma'$  is a restriction of the curve  $\gamma$  to a subinterval or a point.

If  $\alpha: I = [a, b] \rightarrow \mathbb{R}^n$  is a closed curve then its *length* is

$$\ell(\alpha) = \sup \sum_{i=1}^l |\alpha(t_i) - \alpha(t_{i+1})|,$$

where the supremum is taken over all finite partitions  $a = t_1 \leq t_2 \leq \dots \leq t_l \leq t_{l+1} = b$ . If a curve  $\alpha$  is not closed then put its length equal to  $\ell(\alpha) = \sup \ell(\alpha|_J)$ , where the supremum is taken over all closed subintervals  $J$  of  $I$ .

A curve  $\alpha: I \rightarrow \mathbb{R}^n$  is called *rectifiable* whenever  $\ell(\alpha) < \infty$ . A curve is called *locally rectifiable* if each closed subcurve of it is rectifiable.

Consider a closed curve  $\alpha: [a, b] \rightarrow \mathbb{R}^n$  and suppose that it is rectifiable. Define a function  $s_\alpha: [a, b] \rightarrow \mathbb{R}$  by the equality  $s_\alpha(t) = \ell(\alpha|_{[a,t]})$ . For the rectifiable curve  $\alpha$  there exists a unique curve  $\alpha^0: [0, \ell(\alpha)] \rightarrow \mathbb{R}^n$  obtained from  $\alpha$  by a monotonely increasing change of parameter such that  $s_{\alpha^0}(t) = t$  and  $\alpha = \alpha^0 \circ s_\alpha$  [5, Section 2.4]. The curve  $\alpha^0$  is called the *positive natural parametrization* of  $\alpha$ .

Take a Borel set  $A \subset \mathbb{R}^n$  and a Borel function  $\rho: A \rightarrow [0, \infty]$ . The integral of  $\rho$  along a rectifiable curve  $\alpha: [a, b] \rightarrow \mathbb{R}^n$  is defined as

$$\int_{\alpha} \rho ds = \int_0^{\ell(\alpha)} \rho(\alpha^0(\tau)) d\mathcal{H}^1(\tau)$$

with an usual Lebesgue integral in the right-hand side. If  $\alpha$  is absolutely continuous then so is the function  $s_\alpha(t) = [a, b] \rightarrow [0, \ell(\alpha)]$ . Putting  $\tau = s_\alpha(t)$  in the last integral, using the change-of-variables theorem for Lebesgue integrals, and accounting for  $\dot{\alpha}(t) = \frac{d}{dt}\alpha^0(s_\alpha(t))\dot{s}_\alpha(t)$  and  $\frac{d}{d\tau}\alpha^0(\tau) = 1$ , we infer that

$$\int_{\alpha} \rho ds = \int_a^b \rho(\alpha(t)) |\dot{\alpha}(t)| d\mathcal{H}^1(t). \quad (6)$$

Observe that by the change of variable formula we can express this as

$$\int_{\alpha} \rho ds = \int_a^b \rho(\alpha(t)) |\dot{\alpha}(t)| d\mathcal{H}^1(t) = \int_{|\alpha|} \rho(y) \mathcal{N}(y, \alpha, [a, b]) d\mathcal{H}^1(y), \quad (7)$$

where  $\mathcal{N}(y, \alpha, [a, b]) = \#\{[a, b] \cap \alpha^{-1}(y)\}$  is the Banach indicatrix.

For a locally rectifiable curve  $\alpha: I \rightarrow \mathbb{R}^n$ , put

$$\int_{\alpha} \rho ds = \sup_{\beta} \int_{\beta} \rho ds, \quad (8)$$

where the supremum is taken over all closed subcurves  $\beta$  of  $\alpha$ .

Consider a family  $\Gamma$  of curves in  $\mathbb{R}^n$ , where  $n \geq 2$ . A Borel function  $\rho: \mathbb{R}^n \rightarrow [0, \infty]$  is called *admissible* for  $\Gamma$  whenever

$$\int_{\gamma} \rho \, ds \geq 1 \tag{9}$$

for each locally rectifiable curve  $\gamma \in \Gamma$ . Denote the collection of all admissible functions by  $\text{adm } \Gamma$ . Given a weight function  $\theta: \mathbb{R}^n \rightarrow (0, \infty)$  and a number  $p \in [1, \infty)$ , define the  $\theta$ -*weighted*  $p$ -*modulus* of  $\Gamma$  as

$$\text{mod}_p^\theta \Gamma = \inf_{\rho \in \text{adm } \Gamma} \int_{\mathbb{R}^n} \rho^p \theta \, d\mathcal{H}^n.$$

Properties of the weight function will be prescribed separately; at least, we assume that it is locally summable and  $0 < \theta < \infty$  holds  $\mathcal{H}^n$ -almost everywhere. For  $\theta \equiv 1$  we obtain the usual definition of  $p$ -modulus, and instead of  $\text{mod}_p^1 \Gamma$  we write  $\text{mod}_p \Gamma$ . If  $\text{adm } \Gamma = \emptyset$  then we put  $\text{mod}_p^\theta \Gamma = \infty$ ; this case is realized only if  $\Gamma$  contains at least one curve determining a constant mapping.

REMARK 2. The definition of modulus implies that every family of curves which are not locally rectifiable has zero modulus. Moreover, if  $\Gamma$  is a family of curves and  $\Gamma_1 = \{\gamma \in \Gamma : \gamma \text{ is locally rectifiable}\}$  then  $\text{mod}_p^\theta(\Gamma) = \text{mod}_p^\theta(\Gamma_1)$ .

Suppose that  $\alpha$  is a rectifiable closed curve in  $\mathbb{R}^n$ . A mapping  $g: |\alpha| \rightarrow \mathbb{R}^n$  is called *absolutely continuous on  $\alpha$*  if the composition  $g \circ \alpha^0$  is absolutely continuous on  $[0, \ell(\alpha)]$ .

**Theorem 2** [5, Fuglede’s Theorem; 6]. *Suppose that  $f: \Omega \rightarrow \mathbb{R}^n$  is a mapping of class  $W_p^1(\Omega)$  with  $1 \leq p < \infty$ , and  $\Gamma$  is a family of locally rectifiable curves in  $\Omega$  such that each curve has a closed subcurve on which  $f$  is not absolutely continuous. Then  $\text{mod}_p \Gamma = 0$ .*

## 2. Modification of Theorem 1 in the Case of $n = 2$ and $p = \infty$

In this case parameters  $q, p$  may be taken within  $(1, \infty]$ :  $1 < q \leq p \leq \infty$ . The case  $1 < q \leq p < \infty$  is taken into consideration in Theorem 1.

**Theorem 3.** *Let  $1 < q < p = \infty$ . Suppose that  $\Omega \subset \mathbb{R}^2$  is a domain, and  $f: \Omega \rightarrow \mathbb{R}^2$  is a mapping with inner bounded  $\omega$ -weighted  $(q, \infty)$ -codistortion ( $f \in \mathcal{S}\mathcal{D}(\Omega; q, \infty; \omega, 1)^1$ ), while the weight function  $\theta(x) = \omega^{-\frac{1}{q-1}}(x)$  is locally summable. If  $\Gamma$  is a family of curves in the domain  $\Omega$  then we have the inequality*

$$(\text{mod}_1 f(\Gamma)) \leq \mathcal{K}_{q, \infty}^{\omega, 1}(f; \Omega) (\text{mod}_r^\theta \Gamma)^{1/r} \tag{10}$$

with  $r = \frac{q}{q-1}$ .

In this theorem  $\mathcal{K}_{q, \infty}^{\omega, 1}(f; \Omega) = \|\mathcal{K}_{q, \infty}^{\omega, 1}(\cdot, f) \mid L_r(\Omega)\|$ .

**Theorem 4.** *Suppose that  $\Omega \subset \mathbb{R}^2$  is a domain, and  $f: \Omega \rightarrow \mathbb{R}^2$  is a mapping belonging to the Sobolev class  $W_{1, \text{loc}}^1(\Omega)$  with the nonnegative Jacobian determinant:  $\det Df(x) \geq 0$  for almost all  $x \in \Omega$ . Assume that*

- 1)  $f$  is continuous, open and discrete;
- 2) the mapping  $f$  has bounded codistortion:  $\text{adj } Df(x) = 0$  a. e. on the set  $Z = \{x \in \Omega : \det Df(x) = 0\}$ .

---

<sup>1</sup> In the case  $p = \infty$  we have to replace  $\det Df(x)^{\frac{1}{p}}$  in (1) by 1.

Let, for a weight  $\omega: \mathbb{R}^n \rightarrow [0, \infty]$ ,  $(\infty, \infty)$ -codistortion function

$$\Omega \ni x \mapsto \mathcal{K}_{\infty, \infty}^{\omega, 1}(x, f) = \begin{cases} \omega(x) |\operatorname{adj} Df(x)| & \text{if } \det Df(x) > 0, \\ 0 & \text{otherwise,} \end{cases} \quad (11)$$

belongs to  $L_\infty(\Omega)$  (in another words  $f \in \mathcal{SD}(\Omega; \infty, \infty; \omega, 1)$ ). If the weight function  $\theta(x) = \omega^{-1}(x)$  is locally summable then, for any family of curves  $\Gamma$  in the domain  $\Omega$ , we have the inequality

$$\operatorname{mod}_1 f(\Gamma) \leq \mathcal{K}_{\infty, \infty}^{\omega, 1}(f; \Omega) \operatorname{mod}_1^\theta \Gamma. \quad (12)$$

In this theorem  $\mathcal{K}_{\infty, \infty}^{\omega, 1}(f; \Omega) = \|\mathcal{K}_{\infty, \infty}^{\omega, 1}(\cdot, f) \mid L_\infty(\Omega)\|$ .

Theorems 3 and 4 will be proved in Section 6.

### 3. Application

In paper [7, Example 32] the following class of mappings is considered. Suppose that  $n - 1 < p < \infty$ , and consider a continuous, open and discrete mapping  $f: D' \rightarrow \mathbb{R}^n$  of an open connected domain  $D' \subset \mathbb{R}^n$ , where  $n \geq 2$ , such that

- (1)  $f \in W_{n-1, \operatorname{loc}}^1(D')$ ;
- (2)  $\det Df(y) \geq 0$  and  $f$  has finite codistortion; i. e.,  $\operatorname{adj} Df(y) = 0$   $\mathcal{H}^n$ -almost everywhere on  $Z = \{y \in D' : \det Df(y) = 0\}$ ;
- (3) the inner operator distortion function

$$D' \ni y \mapsto \mathcal{K}_{n-1, s}^{1, 1}(y, f) = \begin{cases} \frac{|\operatorname{adj} Df(y)|}{\det Df(y)^{\frac{n-1}{s}}} & \text{if } \det Df(y) \neq 0, \\ 0 & \text{otherwise,} \end{cases} \quad (13)$$

belongs to  $L_{p, \operatorname{loc}}(D')$ , where  $\frac{1}{p} = \frac{n-1}{n-1} - \frac{n-1}{s}$  holds with  $s = \frac{(n-1)p}{p-1} > n - 1$ ;

- (4) the weight function  $\sigma$  defined as

$$\sigma(y) = \begin{cases} \frac{|\operatorname{adj} Df(y)|^p}{\det Df(y)^{p-1}} & \text{if } y \in D' \setminus Z', \\ 1 & \text{otherwise,} \end{cases} \quad (14)$$

is in  $L_{1, \operatorname{loc}}(D')$ , here  $Z' = \{y \in D' : Df(y) = 0\}$ .

Taking into account saying above we see that  $f: D' \rightarrow D$  meets the assumptions of Theorem 1 with  $D'$  instead of  $\Omega$ :

- (2a)  $f \in W_{n-1, \operatorname{loc}}^1(D')$ ;
- (2b)  $\det Df(y) \geq 0$  and  $f$  has finite codistortion;
- (2c)  $f: D' \rightarrow D$  is a mapping of bounded  $\omega$ -weighted  $(s, s)$ -codistortion with  $\omega(y) = \sigma^{-\frac{1}{p-1}}(y)$ , that is, the  $\omega$ -weighted  $(s, s)$ -codistortion function

$$D' \ni y \mapsto \mathcal{K}_{s, s}^{\omega, 1}(y, f) = \begin{cases} \frac{\omega^{\frac{n-1}{s}}(y) |\operatorname{adj} Df(y)|}{\det Df(y)^{\frac{n-1}{s}}} & \text{if } J(y, f) \neq 0, \\ 0 & \text{otherwise,} \end{cases}$$

belongs to  $L_\infty(D')$  and

$$\|\mathcal{K}_{s, s}^{\omega, 1}(\cdot, f) \mid L_\infty(D')\| = 1 \quad (15)$$

(the last equality is proved in [7, Theorem 3] under more general assumption).

Taking into account saying above, by Theorem 1, we come to the following statement.

**Proposition 1.** *Suppose that a continuous, open and discrete mapping  $f : D' \rightarrow \mathbb{R}^n$  of an open connected domain  $D' \subset \mathbb{R}^n$ , where  $n \geq 2$ , has the following properties:*

- (1)  $f \in W_{n-1, \text{loc}}^1(D')$ ;
- (2)  $\det Df(y) \geq 0$  and  $f$  has finite codistortion ( $\text{adj } Df(y) = 0$   $\mathcal{H}^n$ -almost everywhere on  $Z = \{y \in D' : \det Df(y) = 0\}$ );
- (3) the inner operator distortion function

$$D' \ni y \mapsto \mathcal{K}_{n-1, s}^{1,1}(y, f) = \begin{cases} \frac{|\text{adj } Df(y)|}{\det Df(y)^{\frac{n-1}{s}}} & \text{if } \det Df(y) \neq 0, \\ 0 & \text{otherwise,} \end{cases} \quad (16)$$

belongs to  $L_{p, \text{loc}}(D')$  with some  $p > n-1$ , where  $\frac{1}{p} = \frac{n-1}{n-1} - \frac{n-1}{s}$  holds with  $s = \frac{(n-1)p}{p-1} > n-1$ .

If  $\Gamma$  is a family of curves in the domain  $D'$  then we have the inequality

$$\text{mod}_p f(\Gamma) \leq \text{mod}_p^\sigma \Gamma \quad (17)$$

where the weight function  $\sigma$  is defined in (7).

◁ When deriving inequality (17) the properties (2a)–(2c) formulated above, should be taken into account. Really, we see that  $f \in \mathcal{SD}(\Omega; q, p; \omega, 1)$  with  $q = p = s$  and  $\omega(y) = \sigma^{-\frac{1}{p-1}}(y)$ . Therefore, by Theorem 1, we get the inequality

$$(\text{mod}_{s'} f(\Gamma))^{1/s'} \leq \mathcal{K}_{s, s}^{\omega, 1}(f; D') (\text{mod}_{s'}^\theta \Gamma)^{1/s'}$$

with  $s' = \frac{s}{s-(n-1)}$  (here  $\mathcal{K}_{s, s}^{\omega, 1}(f; D') = \|\mathcal{K}_{s, s}^{\omega, 1}(\cdot, f) \mid L_\infty(D')\|$ ). Because of (15),  $s' = p$  and  $\theta(y) = \omega^{-\frac{1}{s-(n-1)}}(y) = \sigma(y)$  inequality (17) holds. ▷

Taking into account [2, Theorem 34] or [4, Theorem 5.2] and its proof we come to

**Proposition 2.** *Suppose that for a continuous, open and discrete mapping  $f : D' \rightarrow \mathbb{R}^n$  of an open connected domain  $D' \subset \mathbb{R}^n$ , where  $n \geq 2$ , conditions of Proposition 1 hold. If  $E = (A, C)$  is a condenser in  $\Omega$ , then the estimate holds:  $\text{cap}_p f(E) \leq \text{cap}_p^\sigma E$ .*

#### 4. The Special Case of the Mappings Under Consideration: $n = 2$

In the case  $n = 2$  we have the following modification of the results of the previous section.

We have  $1 < p < \infty$  and a continuous, open and discrete mapping  $f : D' \rightarrow \mathbb{R}^2$  of on open connected domains  $D' \subset \mathbb{R}^2$  such that

- (1)  $f \in W_{1, \text{loc}}^1(D')$ ;
- (2)  $\det Df(y) \geq 0$  and  $f$  has finite codistortion; i. e.,  $\text{adj } Df(y) = 0$   $\mathcal{H}^2$ -almost everywhere on  $Z = \{y \in D' : \det Df(y) = 0\}$ ;
- (3) the inner operator distortion function

$$D' \ni y \mapsto \mathcal{K}_{1, \frac{p}{p-1}}^{1,1}(y, f) = \begin{cases} \frac{|\text{adj } Df(y)|}{\det Df(y)^{\frac{p-1}{p}}} & \text{if } \det Df(y) \neq 0, \\ 0 & \text{if } \det Df(y) = 0, \end{cases}$$

belongs to  $L_{p, \text{loc}}(D')$ .

(4) the weight function  $\sigma$  defined as

$$\sigma(y) = \begin{cases} \frac{|\operatorname{adj} Df(y)|^p}{\det Df(y)^{p-1}} & \text{if } y \in D' \setminus Z', \\ 1 & \text{otherwise,} \end{cases} \quad (18)$$

is in  $L_{1,\text{loc}}(D')$ , here  $Z' = \{y \in D' : Df(y) = 0\}$ .

It is not hard to see that the continuous, open and discrete mapping  $f : D' \rightarrow \mathbb{R}^2$  meets the assumptions of Proposition 1 under  $n = 2$ :

(3a)  $f \in W_{1,\text{loc}}^1(D')$ ;

(3b)  $f$  has finite distortion;

(3c)  $f : D' \rightarrow D$  is a mapping with bounded  $\omega$ -weighted  $(p', p')$ -distortion where  $p' = \frac{p}{p-1}$  and  $\omega(y) = \sigma^{-\frac{1}{p-1}}(y)$ , that is the  $\omega$ -weighted  $(p', p')$ -distortion function

$$D' \ni y \mapsto K_{p',p'}^{\omega,1}(y, f) = \begin{cases} \frac{\omega^{\frac{1}{p'}}(y) |Df(y)|}{\det Df(y)^{\frac{1}{p'}}} & \text{if } \det Df(y) \neq 0, \\ 0 & \text{otherwise,} \end{cases}$$

belongs to  $L_\infty(D')$ , and

$$\|K_{p',p'}^{\omega,1}(\cdot, f) | L_\infty(D')\| = 1. \quad (19)$$

Taking into account saying above, by Proposition 1, we come to the following statement.

**Corollary 1.** *Suppose that a continuous, open and discrete mapping  $f : D' \rightarrow \mathbb{R}^2$  of an open connected domain  $D' \subset \mathbb{R}^2$  has the following properties:*

(1)  $f \in W_{1,\text{loc}}^1(D')$ ;

(2)  $f$  has finite codistortion ( $\operatorname{adj} Df(y) = 0$   $\mathcal{H}^2$ -almost everywhere on  $Z = \{y \in D' : \det Df(y) = 0\}$ );

(3) the inner operator distortion function

$$D' \ni y \mapsto \mathcal{K}_{1,p'}^{1,1}(y, f) = \begin{cases} \frac{|\operatorname{adj} Df(y)|}{\det Df(y)^{\frac{1}{p'}}} & \text{if } \det Df(y) \neq 0, \\ 0 & \text{otherwise,} \end{cases} \quad (20)$$

belongs to  $L_{p,\text{loc}}(D')$  with some  $p > 1$ , where  $\frac{1}{p} + \frac{1}{p'} = 1$ .

If  $\Gamma$  is a family of curves in the domain  $D'$  then we have the inequality

$$\operatorname{mod}_p f(\Gamma) \leq \operatorname{mod}_p^\sigma \Gamma \quad (21)$$

holds where the weight function  $\sigma$  is defined in (18).

## 5. One More Special Case of the Mappings Under Consideration: $n = 2$ and $p = 1$

In this section we prove that Corollary 1 is valid also in the case  $p = 1$ . To show this we have to modify some arguments of the previous section. A counterpart of Corollary 1 is formulated in the following statement.

**Proposition 3.** *Suppose that a continuous, open and discrete mapping  $f : D' \rightarrow \mathbb{R}^2$  of an open connected domain  $D' \subset \mathbb{R}^2$  has the following properties:*

(1)  $f \in W_{1,\text{loc}}^1(D')$ ;

(2)  $\det Df(y) \geq 0$  and  $f$  has finite codistortion ( $\operatorname{adj} Df(y) = 0$   $\mathcal{H}^2$ -almost everywhere on  $Z = \{y \in D' \mid \det Df(y) = 0\}$ );

(3) the inner operator codistortion function

$$D' \ni y \mapsto \mathcal{K}_{1,\infty}^{1,1}(y, f) = \begin{cases} |\operatorname{adj} Df(y)| & \text{if } \det Df(y) \neq 0, \\ 0 & \text{otherwise,} \end{cases} \quad (22)$$

belongs to  $L_{1,\operatorname{loc}}(D')$ .

If  $\Gamma$  is a family of curves in  $D'$  then we have

$$\operatorname{mod}_1 f(\Gamma) \leq \operatorname{mod}_1^\sigma \Gamma \quad (23)$$

with  $\sigma$  defined in (25).

◁ We show that the proof of Proposition 3 can be reduced to Theorem 3. For doing this formulate first additional properties of  $f$  and  $\varphi = f^{-1}$ .

PROPERTIES OF  $\varphi = f^{-1}$ . If  $f : D' \rightarrow D$  is a homeomorphism then the inverse homeomorphism  $\varphi = f^{-1} : D \rightarrow D'$  enjoys the following properties:

(4) by [9, Theorem 4] or [7, Theorem 27] we have  $\varphi \in W_{1,\operatorname{loc}}^1(D)$  (see also [10, Theorem 3.2]);

(5)  $\varphi$  has finite distortion by [7, Theorem 27] (see also [10, Theorem 3.3]);

(6)  $\varphi$  is differentiable a. e. in  $D$  by [7, Theorem 27];

while  $f : D' \rightarrow D$

(6)  $\varphi$  belongs to  $\mathcal{Q}_{1,1}(D, D'; \sigma)$  (see [4]), that is the distortion function

$$D \ni x \mapsto K_{1,1}^{1,\sigma}(x, \varphi) = \begin{cases} \frac{|D\varphi(x)|}{\sigma(\varphi(x)) \det D\varphi(x)} & \text{if } \det D\varphi(x) \neq 0, \\ 0 & \text{if } \det D\varphi(x) = 0, \end{cases} \quad (24)$$

of the inverse mapping  $\varphi = f^{-1}$  with the weight function  $\sigma \in L_{1,\operatorname{loc}}(D')$  defined as

$$\sigma(y) = \begin{cases} |\operatorname{adj} Df(y)| & \text{if } y \in D' \setminus Z', \\ 1 & \text{otherwise,} \end{cases} \quad \text{where } Z' = \{y \in D' : Df(y) = 0\}, \quad (25)$$

is in  $L_\infty(D)$  and  $K_{1,1}^{1,\sigma}(\varphi; D) = \|K_{1,1}^{1,\sigma}(\cdot, \varphi) \mid L_\infty(D)\| = 1$  (see [4, Theorem 25, formulas (30) and (37); 8]).

PROPERTIES OF  $f$ . Taking into account saying above, we see that  $f : D' \rightarrow D$  meets some additional properties:

(7)  $f \in W_{1,\operatorname{loc}}^1(D')$  and  $f$  is differentiable a. e. in  $D'$  by [7, Theorem 27];

(8)  $\det Df(y) \geq 0$  and  $f$  has finite distortion by [7, Theorem 27] (see also [10, Theorem 3.3]);

(9)  $f : D' \rightarrow D$  is a mapping with bounded  $\omega$ -weighted  $(\infty, \infty)$ -codistortion with the weight function  $\omega = \sigma^{-1}$ , that is the  $\omega$ -weighted  $(\infty, \infty)$ -codistortion function

$$D' \ni y \mapsto \mathcal{K}_{\infty,\infty}^{\omega,1}(y, f) = \begin{cases} \omega(y) |\operatorname{adj} Df(y)| & \text{if } \det Df(y) \neq 0, \\ 0 & \text{otherwise,} \end{cases}$$

belongs to  $L_\infty(D')$ , and

$$\|\mathcal{K}_{\infty,\infty}^{\omega,1}(\cdot, f) \mid L_\infty(D')\| = \|K_{1,1}^{1,\sigma}(\cdot, \varphi) \mid L_\infty(D)\| = 1. \quad (26)$$

Now it is evident that  $f$  enjoys the conditions of Theorem 3, and therefore (23) holds for  $f$ . ▷

## 6. Proof of Theorems 3 and 4

◁ We verify that the proof of Theorem 1 given in [4, Theorem 4.1] for mappings with bounded  $\theta$ -weighted  $(q, p)$ -codistortion, where  $n - 1 < q \leq p < \infty$ , works also in the case  $1 < q \leq p = \infty$  at  $n = 2$ . To do this we need properties of Poletsky function and Poletsky's Lemma in this case. We formulate and prove them below. ▷

**1. Properties of Poletsky function.** Take a continuous mapping  $f : \Omega \rightarrow \mathbb{R}^2$  and a domain  $D$  compactly embedded into  $\Omega$ , meaning that  $D$  is bounded and  $\bar{D} \subset \Omega$ , written briefly as  $D \Subset \Omega$ , and take  $y \notin f(\partial D)$ . Denote by  $\mu(y, f, D)$  the degree of  $f$  at  $y$  with respect to  $D$ . Say that  $f$  is sense-preserving whenever  $\mu(y, f, D) > 0$  for all domains  $D \Subset \Omega$  and all points  $y \in f(D) \setminus f(\partial D)$ . For  $A \subset \Omega$  refer as the multiplicity function to  $\mathbb{R}^2 \ni y \mapsto N(y, f, A) = \# \{f^{-1}(y) \cap A\}$ . Moreover, put  $N(f, A) = \sup_{y \in \mathbb{R}^2} N(y, f, A)$ .

Suppose that  $f : \Omega \rightarrow \mathbb{R}^2$  is a continuous, open, and discrete mapping. A domain  $D \Subset \Omega$  is called normal whenever  $f(\partial D) = \partial f(D)$ . A normal neighborhood of  $x \in \Omega$  is a normal domain  $U \subset \Omega$  such that  $\bar{U} \cap f^{-1}(f(x)) = \{x\}$ . The quantity  $i(x, f) = \mu(f(x), f, U)$  is independent of the choice of a normal neighborhood  $U$  of  $x$  (see [11, Chapter II, §2] for instance) and is called the local index of  $f$  at  $x$ . A point  $x \in \Omega$  is called a branch point of  $f$  whenever  $f$  is not a homeomorphism of any neighborhood of  $x$ . Denote the collection of all branch points of  $f$  by  $B_f$ . If  $D$  is a normal domain for a mapping  $f$  then  $\mu(y, f, D)$  is independent of  $y \in f(D)$ . We will call this constant by  $\mu(f, D)$ .

In the following two lemmas we state propositions of interest in their own right. Both of them are applied in the proof of the main result of this section.

**Lemma 1** [3, Lemma 10]. *Assume that  $f : \Omega \rightarrow \mathbb{R}^2$  is a continuous, open and discrete mapping in  $W_{1,\text{loc}}^1(\Omega)$  with finite distortion. Then for every open connected set  $U \subset \Omega$  the set  $\{x \in U \setminus B_f : J(x, f) \neq 0\}$  has positive measure.*

◁ If, on the contrary,  $J(x, f) = 0$  a.e. on a connected set  $U \subset \Omega \setminus B_f$  on which  $f$  is a homeomorphism then  $Df(x) = 0$  a.e. on  $U$  because  $f$  has finite distortion. Then  $f$  is constant on  $U$ , and consequently,  $f$  cannot be open. ▷

**Proposition 4.** *If  $f : \Omega \rightarrow \mathbb{R}^2$  is a continuous, open and discrete mapping in  $W_{1,\text{loc}}^1(\Omega)$  with finite distortion, then  $f$  is differentiable a.e. on  $\Omega \setminus B_f$  and sense-preserving.*

◁ For a connected open set  $U \subset \Omega \setminus B_f$  on which  $f$  is a homeomorphism, it is enough to apply the statement [9, Theorem 4] or [7, Theorem 27] twice. For the restriction  $f|_U : U \rightarrow f(U)$  it provides that the inverse homeomorphism  $(f|_U)^{-1} : f(U) \rightarrow U$  is in  $W_1^1(f(U))$ , is of finite distortion, and is differentiable a.e. on  $f(U)$ . Then applying [7, Theorem 27] to  $(f|_U)^{-1} : f(U) \rightarrow U$  we get similar properties to the given mapping  $f|_U : U \rightarrow f(U)$ . By Lemma 1,  $\det Df(x) \geq 0$  and properties of degree we conclude that  $f$  is sense-preserving. ▷

**DEFINITION 3.** For a sense-preserving, continuous, open and discrete mapping  $f : \Omega \rightarrow \mathbb{R}^2$  and a normal domain  $D \Subset \Omega$ , define the Poletsky function  $g_D : V \rightarrow \mathbb{R}^2$  on  $V = f(D)$  [12] by putting

$$V \ni y \mapsto g_D(y) = \Lambda \sum_{x \in f^{-1}(y) \cap D} i(x, f)x, \quad (27)$$

where  $\Lambda = \mu(f, D)$ .

The function of the form (27) was introduced by Poletsky in [12] for mappings with bounded distortion ( $p = q = n$ ,  $\omega \equiv 1$ ). The next statement presents the properties of the Poletsky function for the classes of mappings under consideration.

**Proposition 5** [2, 3]. Suppose that  $f : \Omega \rightarrow \mathbb{R}^2$  belongs to  $\mathcal{O}\mathcal{D}(\Omega; \infty, \infty; \omega, 1)$  (properties (4a)–(4c) hold). Then

- (1) the function  $g_D$  defined in (27) is continuous and belongs to  $\text{ACL}(V)$ ;
- (2)  $Dg_D(y) = 0$  a. e. on  $Z' \cup \Sigma'$ ;
- (3) Poletsky function  $g_D$  defined in (27) is in  $W_1^1(V)$ ; furthermore,

$$\|Dg_D \mid L_1(V)\| \leq \Lambda \|K_{\infty, \infty}^{\omega, 1}(\cdot; f) \mid L_\infty(D)\| \int_D \sigma(x) dx.$$

We emphasize that the formulated statement is proved in [2, Theorem 18] for mappings  $f \in \mathcal{S}\mathcal{D}(\Omega; p, p; \omega, 1)$ ,  $p \in (1, \infty)$ . The same proof works also in the case  $p = \infty$  at  $n = 2$ .

**2. Poletsky’s Lemma.** Consider a continuous, open and discrete mapping  $f : \Omega \rightarrow \mathbb{R}^2$ . Take a closed rectifiable curve  $\beta : I_0 \rightarrow \mathbb{R}^n$  and a curve  $\alpha : I \rightarrow \Omega$  with  $f \circ \alpha \subset \beta$ . In particular, we have  $I \subset I_0$ . If the function  $s_\beta : I_0 \rightarrow [0, \ell(\beta)]$  is constant on some interval  $J \subset I$ , then the mapping  $\beta$  is constant on  $J$ . In turn, since  $f$  is discrete,  $\alpha$  is also constant on  $J$ . Consequently, there exists a unique mapping  $\alpha^* : s_\beta(I) \rightarrow \Omega$  satisfying  $\alpha = \alpha^* \circ s_\beta|_I$ . We can prove that  $\alpha^*$  is continuous and  $f \circ \alpha^* \subset \beta^0$ . The curve  $\alpha^*$  is called an  $f$ -representative of  $\alpha$  (with respect to  $\beta$ ) whenever  $\beta = f \circ \alpha$ . Suppose now that  $\beta = f \circ \alpha$ . The above arguments show that

$$f \circ \alpha^* = (f \circ \alpha)^0.$$

Therefore, the curve  $f \circ \alpha^*$  admits a positive natural parametrization, and hence it is Lipschitz. Thus we can integrate along this curve using (6) where  $|\frac{d}{dt}(f \circ \alpha^*)(t)| = 1$  for  $\mathcal{H}^1$ -almost all  $t \in I$ .

The mapping  $f$  is called *absolutely precontinuous* on  $\alpha$  provided that  $\alpha^*$  is absolutely continuous.

**Lemma 2.** Suppose that  $f : \Omega \rightarrow \mathbb{R}^2$  is a mapping of class  $\mathcal{S}\mathcal{D}(\Omega; \infty, \infty; \omega, 1)$ . Consider a family  $\Gamma$  of curves in  $\Omega$  such that for every  $\gamma \in \Gamma$  the following holds: the curve  $f \circ \gamma$  is locally rectifiable and  $\gamma$  has a closed subcurve  $\alpha$  on which  $f$  is not absolutely precontinuous. Then  $\text{mod}_1 f(\Gamma) = 0$ .

The formulated Lemma is proved in [4, Lemma 3.3] for mappings  $f \in \mathcal{S}\mathcal{D}(\Omega; p, p; \omega, 1)$ ,  $p \in (1, \infty)$ . The same proof works also in the case  $p = \infty$  at  $n = 2$ .

In the proof of Lemma 2 we also need the following statement.

**Lemma 3.** Consider a homeomorphism  $\varphi : \Omega \rightarrow \Omega'$  of class  $\mathcal{S}\mathcal{D}(\Omega; q, \infty; \theta, 1)$ , where  $\Omega, \Omega' \subset \mathbb{R}^2$  and  $1 < q \leq \infty$ .

Then

- (1) the inverse homeomorphism is  $\varphi^{-1} \in W_{1, \text{loc}}^1(\Omega')$ ;
- (2)  $\varphi^{-1}$  has finite distortion:  $D\varphi^{-1}(y) = 0$  almost everywhere on  $Z'$ ;
- (3)  $K_{1, r}^{1, \omega}(\cdot, \varphi^{-1}) \in L_\varrho(\Omega')$ , where

$$r = \begin{cases} \frac{q}{q-n+1} & \text{if } q < \infty, \\ 1 & \text{if } q = \infty, \end{cases} \quad \omega = \begin{cases} \theta^{-\frac{1}{q-1}} & \text{if } q < \infty, \\ \theta^{-1} & \text{if } q = \infty; \end{cases}$$

(4) if the weight function  $\omega$  is locally summable then the inverse homeomorphism induces, by the change-of-variable rule, the bounded operator

$$\varphi^{-1*} : L_r^1(\Omega; \omega) \cap W_{\infty, \text{loc}}^1 \rightarrow L_1^1(\Omega').$$

We have the relations

$$\|K_{1, r}^{1, \omega}(\cdot, \varphi^{-1}) \mid L_\varrho(\Omega')\| = \|\mathcal{K}_{q, \infty}^{\theta, 1}(\cdot, \varphi) \mid L_\varrho(\Omega)\|$$

and

$$\beta_{q,\infty} \|K_{1,r}^{1,\omega}(\cdot, \varphi^{-1}) | L_\rho(\Omega')\| \leq \|\varphi^{-1*}\| \leq \|K_{1,r}^{1,\omega}(\cdot, \varphi^{-1}) | L_\rho(\Omega')\|,$$

where  $\beta_{q,\infty}$  is some constant.

◁ Properties (1) and (2) of  $\varphi = f^{-1}$  were proved just after Proposition 3. Taking into account (1) and (2) Properties (3) and (4) can be proved by analogy with Theorem 9 of [2]. ▷

REMARK 3. By means of Theorems 3 and 4 for homeomorphisms  $\varphi : \Omega \rightarrow \Omega'$  of class  $\mathcal{S}\mathcal{D}(\Omega; q, \infty; \theta, 1)$ , where  $\Omega, \Omega' \subset \mathbb{R}^2$  and  $1 < q \leq \infty$ , we can prove some more inequalities such that Väisälä inequality and the capacity inequality (see proofs in [4, Theorem 22] and [4, Theorem 28] respectively).

REMARK 4. It is not hard to see that assumptions of Theorem 4 are weaker comparing with those in paper [13]. For instance, Theorem 1.3 of [13] is formulated under addition condition that the given mapping is closed. Therefore Theorem 4 with weaker assumptions contains the main result of paper [13].

**Acknowledgements.** I greatly appreciate the anonymous reviewers for critically reading and comments, which helped improve the initial manuscript.

## References

1. Reshetnyak Yu. G. *Space Mappings with Bounded Distortion*, Providence, Amer. Math. Soc., 1989.
2. Vodopyanov, S. K. Basics of the Quasiconformal Analysis of a Two-Index Scale of Space Mappings, *Siberian Mathematical Journal*, 2018, vol. 59, no. 5, pp. 805–834. DOI: 10.1134/S0037446618050075.
3. Vodopyanov, S. K. Differentiability of Mappings of the Sobolev Space  $W_{n-1}^1$  with Conditions on the Distortion Function, *Siberian Mathematical Journal*, 2018, vol. 59, no. 6, pp. 983–1005. DOI: 10.1134/S0037446618060034.
4. Vodopyanov, S. K. Moduli Inequalities for  $W_{n-1,loc}^1$ -Mappings with Weighted Bounded  $(q, p)$ -Distortion, *Complex Variables and Elliptic Equations*, 2021, vol. 66, no. 6–7, pp. 1037–1072. DOI: 10.1080/17476933.2020.1825396.
5. Väisälä J. *Lectures on  $n$ -Dimensional Quasiconformal Mappings*, Lecture Notes in Mathematics, vol. 229, Berlin-Heidelberg-New York, Springer, 1971.
6. Fuglede, B. Extremal Length and Functional Completion, *Acta Mathematica*, 1957, vol. 98, pp. 171–219. DOI: 10.1007/BF02404474.
7. Vodopyanov, S. K. The Regularity of Inverses to Sobolev Mappings and the Theory of  $Q_{q,p}$ -Homeomorphisms, *Siberian Mathematical Journal*, 2020, vol. 61, no. 6, pp. 1002–1038. DOI: 10.1134/S0037446620060051.
8. Vodopyanov, S. K. and Tomilov, A. O. Functional and Analytic Properties of a Class of Mappings in Quasi-Conformal Analysis, *Izvestiya: Mathematics*, 2021, vol. 85, no. 5, pp. 883–931. DOI: 10.1070/IM9082.
9. Vodopyanov, S. K. Regularity of Mappings Inverse to Sobolev Mappings, *Sbornik: Mathematics*, 2012, vol. 203, no. 10, pp. 1383–1410. DOI: 10.1070/SM2012v203n10ABEH004269.
10. Hencl, S. and Koskela, P. Regularity of the Inverse of a Planar Sobolev Homeomorphism, *Archive for Rational Mechanics and Analysis*, 2006, vol. 180, pp. 75–95. DOI: 10.1007/s00205-005-0394-1.
11. Rickman S. *Quasiregular mappings*, Berlin, Springer-Verlag, 1993, 213 p.
12. Poletsky, E. A. The Modulus Method for Nonhomeomorphic Quasiconformal Mappings, *Mathematics of the USSR-Sbornik*, 1970, vol. 12, no. 2, pp. 260–270. DOI: 10.1070/SM1970v012n02ABEH000921.
13. Salimov, R. R., Sevost'yanov, E. A. and Targonskii, V. A. On Modulus Inequality of the Order  $p$  for the Inner Dilatation. *arXiv - MATH - Complex Variables*. 2022. DOI:arxiv-2204.07870.

Received September 2, 2022

SERGEY K. VODOPYANOV  
 Sobolev Institute of Mathematics,  
 4 Akademika Koptyuga Ave., Novosibirsk 630090, Russia,  
 Principal Researcher  
 E-mail: vodopis@math.nsc.ru  
<https://orcid.org/0000-0003-1238-4956>

О МОДУЛЬНЫХ НЕРАВЕНСТВАХ ТИПА ПОЛЕЦКОГО  
ДЛЯ НЕКОТОРЫХ КЛАССОВ ОТОБРАЖЕНИЙВодопьянов С. К.<sup>1</sup><sup>1</sup> Институт математики им. С. Л. Соболева,  
Россия, 630090, Новосибирск, пр-т Академика Коптюга, 4  
E-mail: vodopis@math.nsc.ru

**Аннотация.** Хорошо известно, что теория отображений с ограниченным искажением была заложена Ю. Г. Решетняком в 60-е годы прошлого века [1]. В работах [2, 3] была введена двухиндексная шкала отображений с весовым ограниченным  $(q, p)$ -искажением. Эта шкала отображений включает в себя, в частности, отображения с ограниченным искажением, упомянутые выше (при  $q = p = n$  и тривиальной весовой функции). В работе [4] для двухиндексной шкалы отображений с весовым ограниченным  $(q, p)$ -искажением доказано модульное неравенство типа Полецкого при минимальной регулярности; приведено много примеров отображений, к которым можно применить результаты [4]. В этой статье мы приведем одно такое применение. Другая цель этой статьи — показать новый класс отображений, в которых выполняются модульные неравенства типа Полецкого. Для этого мы расширим при  $n = 2$  справедливость утверждений работы [4] на предельные показатели:  $1 < q \leq p \leq \infty$ . Это обобщение содержит в качестве частного случая результаты недавно опубликованных работ. Как следствие результатов этой статьи мы получаем также оценки изменения емкости конденсаторов.

**Ключевые слова:** квазиконформный анализ, пространство Соболева, модуль семейства кривых, оценка модуля.

**AMS Subject Classification:** 30C65 (26B35, 31B15, 46E35).

**Образец цитирования:** Vodopyanov S. K. On Poletsky-Type Modulus Inequalities for Some Classes of Mappings // Владикавк. мат. журн.—2022.—Т. 24, № 4.—С. 58–69 (in English). DOI: 10.46698/w5793-5981-8894-о.